

## Development of the Eat, Sleep, Console (ESC) Care Tool for Care of Opioid-exposed Newborns

### REFERENCES – Whalen

- 1) Winkelman TNA, Villapiano N, Kozhimannil KB, et al. Incidence and costs of neonatal abstinence syndrome among infants with Medicaid: 2004-2014. *Pediatrics*. 2018;141:e20173520.
- 2) Villapiano NLG, Winkelman TNA, Kozhimannil KB, et al. Rural and urban differences in neonatal abstinence syndrome and maternal opioid use, 2004 to 2013. *JAMA Pediatr*. 2017;171:194-196.
- 3) Milliren CE, Gupta M, Graham DA, et al. Hospital variation in neonatal abstinence syndrome incidence, treatment modalities, resource use, and costs across pediatric hospitals in the United States, 2013 to 2016. *Hosp Pediatr*. 2018;8:15-20.
- 4) Patrick SW, Davis MM, Lehmann CU, Cooper WO. Increasing incidence and geographic distribution of neonatal abstinence syndrome: United States 2009 to 2012. *J Perinatol*. 2015;35:650–655.
- 5) Haight SC, Ko JK, Tong VT, et al. Opioid use disorder documented at delivery hospitalization - United States, 1999-2014. *MMWR*. 2018;67:845-849.
- 6) Kocherlakota P. Neonatal abstinence syndrome. *Pediatrics*. 2014;134:e547–561.
- 7) Mehta A, Forbes KD, Kuppala VS. Neonatal abstinence syndrome management from prenatal counseling to postdischarge follow-up care: results of a national survey. *Hosp Pediatr*. 2013;3:317-323.
- 8) Bogen DL, Whalen BL, Kair LR, et al. Wide variation found in care of opioid-exposed newborns. *Acad Pediatr*. 2017;17:374-380.
- 9) Murphy K, Coo H, Warre R, et al. Variations and similarities in clinical management of neonatal abstinence syndrome: Findings of a Canadian survey. *Paediatr Child Health*. 2017;22:148-152.
- 10) Snowden JN, Akshatha A, Annett RD, et al. The ACT NOW clinical practice survey: gaps in the care of infants with neonatal opioid withdrawal syndrome. *Hosp Pediatr*. 2019;9:585–592.
- 11) Finnegan LP, Connaughton JF Jr, Kron RE, Emich JP. Neonatal abstinence syndrome: assessment and management. *Addict Dis*. 1975;2:141–158.

- 12) Finnegan LP, Kron RE, Connaughton JF, Emich JP. Assessment and treatment of abstinence in the infant of the drug-dependent mother. *Int J Clin Pharmacol Biopharm.* 1975;12:19-32.
- 13) D'Apolito K, Finnegan L. Assessing signs & symptoms of neonatal abstinence using the Finnegan scoring tool. An inter-observer reliability program. Second ed. Instructional Manual. © *Neo Advances LLC*; 2010.
- 14) Murphy K, Coo H, Warre R, et al. Variations and similarities in clinical management of neonatal abstinence syndrome: Findings of a Canadian survey. *Paediatr Child Health.* 2017;22:148-152.
- 15) Walsh MC, Crowley M, Wexelblatt S, et al. Ohio perinatal quality collaborative improves care of neonatal narcotic abstinence syndrome. *Pediatrics.* 2018;141 (4) e20170900.
- 16) Kraft WK, Gibson E, Dysart K. Sublingual buprenorphine for treatment of neonatal abstinence syndrome: a randomized trial. *Pediatrics.* 2008;122:e601-617.
- 17) Patrick SW, Kaplan HC, Passarella M, et al. Variation in treatment of neonatal abstinence syndrome in US Children's Hospitals, 2004-2011. *J Perinatol.* 2014;34:867–872.
- 18) Hall ES, Wexelblatt SL, Crowley M, et al. A multicenter cohort study of treatments and hospital outcomes in neonatal abstinence syndrome. *Pediatrics.* 2014;134:e527–e534.
- 19) Hall ES, Isemann BT, Wexelblatt SL, et al. A cohort comparison of buprenorphine versus methadone treatment for neonatal abstinence Syndrome. *J Pediatr,* 2016;170:39–44.
- 20) Kraft WK, Adeniyi-Jones SC, Chervoneva I, et al. Buprenorphine for the treatment of the neonatal abstinence syndrome. *N Engl J Med.* 2017;376:2341–2348.
- 21) Tolia VN, Patrick SW, Bennett MM, et al. Increasing incidence of the neonatal abstinence syndrome in U.S. neonatal ICUs. *N Engl J Med.* 2015;372:2118–2126.
- 22) Davis JM, Shenberger J, Terrin N, et al. Comparison of Safety and Efficacy of Methadone vs. Morphine for Treatment of Neonatal Abstinence Syndrome: A Randomized Clinical Trial. *JAMA Pediatr.* 2018;172:741–748.
- 23) Hall ES, Rice WR, Folger AT, Wexelblatt SL. Comparison of Neonatal Abstinence Syndrome Treatment with Sublingual Buprenorphine versus Conventional Opioids. *Am J Perinatol.* 2018;35:405-412.
- 24) Zimmermann-Baer U, Notzli U, Rentsch K, Bucher HU. Finnegan neonatal abstinence scoring system: normal values for first 3 days and weeks 5–6 in non-addicted infants. *Addiction.* 2010;105:524–528.
- 25) D'Apolito K. Assessing Neonates for Neonatal Abstinence: Are you reliable? *J Perinat Neonatal Nurs.* 2014;28:220-231.

- 26) Retskin CM and Wright ME. Interobserver Reliability of the Finnegan Neonatal Abstinence Scoring Tool in an Acute Care Setting. *JOGNN*. 2014;43:S61.
- 27) Asti L, Magers JS, Keels E, et al. A quality improvement project to reduce length of stay for neonatal abstinence syndrome. *Pediatrics*. 2015;135:e1494-1500.
- 28) Patrick SW, Schumacher Re, Horbar JD, et al. Improving care for neonatal abstinence syndrome. *Pediatrics*. 2016;137 (5) e20153835.
- 29) Atwood EC, Sollender G, Hsu E, et al. A qualitative study of family experience with hospitalization for neonatal abstinence syndrome. *Hosp Pediatr*. 2016;6:626-632.
- 30) Cleveland LM, Bonugli R. Experiences of mothers of infants with neonatal abstinence syndrome in the neonatal intensive care unit. *J Obstet Gynecol Neonatal Nurs*. 2014;43:318-329.
- 31) Zahorodny W, Rom C, Whitney W, Giddens S, Samuel M, Maichuk G, Marshall R. The neonatal withdrawal inventory: a simplified score of newborn withdrawal. *J Dev Behav Pediatr*. 1998;19:89-93.
- 32) American Academy of Pediatrics Committee on Drugs. Neonatal drug withdrawal. *Pediatrics*. 1998;101:1079-1088. [published correction appears in *Pediatrics*. 1998;201:660].
- 33) Maguire D, Cline GJ, Parnell L, Tai CY. Validation of the Finnegan neonatal abstinence syndrome tool – short form. *Adv Neonatal Care*. 2013;13:430-437.
- 34) Jones HE, Seashore C, Johnson E, Horton E, O’Grady KE, Andringa K. Measurement of neonatal abstinence syndrome: Evaluation of short forms. *J Opioid Manag*. 2016;12:19-23.
- 35) Gomez Pomar E, Finnegan LP, Devlin L, Bada H, Concina VA, Ibonia KT, Westgate PM. Simplification of the Finnegan Neonatal Abstinence Scoring System: retrospective study of two institutions in the USA. *BMJ Open*. 2017;7:e016176.
- 36) Chervoneva I, Adeniyi-Jones SC, Blanco F, Kraft WK. Development of an abbreviated symptom score for the neonatal abstinence syndrome. *J Perinatol*. 2020;40:1031-1040.
- 37) Devlin LA, Breeze JL, Terrin N, Gomez Pomar E, Bada H, Finnegan LP, O’Grady KE, Jones HE, Lester B, Davis JM. Association of a Simplified Finnegan Neonatal Abstinence Scoring Tool With the Need for Pharmacologic Treatment for Neonatal Abstinence Syndrome. *JAMA Netw Open*. 2020;3:e202275.
- 38) Kocherlakota P, Qian EC, Patel VC, Mandru C, Vilar RE, Alpan G, La Gamma EF. A New Scoring System for the Assessment of Neonatal Abstinence Syndrome. *Am J Perinatol*. 2020;37:333-340.
- 39) Holmes AV, Atwood EC, Whalen B, et al. Rooming-In to Treat Neonatal Abstinence Syndrome: Improved Family-Centered Care at Lower Cost. *Pediatrics*. 2016;137:e20152929.
- 40) Wachman EM, Grossman M, Schiff DM, et al. Quality improvement initiative to improve inpatient outcomes for Neonatal Abstinence Syndrome. *J Perinatol*. 2018;38:1114-1122.

- 41) Dr. Matt Grossman's National Center on Substance Abuse and Child Welfare (NCSACW) 3/21/18 webinar "Infants with Prenatal Substance Exposure and their Parents: Family Approach of Yale New Haven Children's Hospital"  
<https://www.youtube.com/watch?v=7epcyi2mafY>.
- 42) Grossman MR, Berkwitz AK, Osborn RR, et al. An Initiative to Improve the Quality of Care of Infants with Neonatal Abstinence Syndrome. *Pediatrics*. 2017;139:e20164460.
- 43) Grossman MR, Lipshaw MJ, Osborn RR, Berkwitz AK. A Novel Approach to Assessing Infants with Neonatal Abstinence Syndrome. *Hosp Pediatr*. 2018;8:1-6.
- 44) MacMillan KDL, Whalen BL, Flanagan V, et al. Understanding Family Perspectives: A Follow-Up Qualitative Study of Family Experience with Hospitalization for Neonatal Abstinence Syndrome. *Pediatric Academic Societies' Annual Meeting*. May 2018.
- 45) Wachman EM, Houghton M, Melvin P, et al. A quality improvement initiative to implement the eat, sleep, console neonatal opioid withdrawal syndrome care tool in Massachusetts' PNQIN collaborative. *J Perinatol*. 2020;1-10.